

- Kampo product -

JUNKOU Goshuyuto FC Extract Fine Granules for Ethical Use
(Goshuyuto)

Storage: Store at room temperature.
See the section "PRECAUTIONS FOR HANDLING"
Expiration date: The expiration date is indicated on the
outer package.

Approval No.	(61AMY) 0378
Date of listing in the NHI reimbursement price	October 1986
Date of initial marketing in Japan	October 1986

DESCRIPTION

(1) The daily dose of this product, 6.00 g, contains 2.80 g of the dried extract (Goshuyuto extract) from the following mixed crude drugs.

JP Ginseng ----- 3.00 g

JP Ginger ----- 1.50 g

JP Jujube ----- 3.00 g

JP Euodia Fruit ----- 4.00 g

(JP: The Japanese Pharmacopeia)

The inactive ingredients contained are Corn Starch and Lactose Hydrate.

(2) This product is light brown-colored fine granules, smells uniquely, and tastes hot initially and lingeringly bitter later.
ID Code: FC 31

INDICATIONS

The following symptoms of those patients with distended stomach pit who have cold hands and feet:
Headache, nausea accompanying headache, and hiccup

DOSAGE AND ADMINISTRATION

The usual adult dose is 6.0 g/day orally in 2 or 3 times before or between meals. The dosage may be adjusted according to the patient's age, body weight, and symptoms.

PRECAUTIONS**(1) Important Precautions**

- When this product is used, the patient's "SHO" (constitution/symptoms) should be taken into account. The patient's progress should be carefully monitored, and if no improvement in symptoms/findings is observed, continuous treatment should be avoided.
- When this product is coadministered with other Kampo-products (Japanese traditional herbal medicines), etc., attention should be paid to the duplication of the contained crude drugs.

(2) Adverse Reactions

This product has not been investigated (drug use investigations, etc.) to determine the incidence of adverse reactions. Therefore, the incidence of adverse reactions is not known.

	Incidence Unknown
Hypersensitivity Note 1)	Rash, Urticaria, etc.
Hepatic	Abnormality of hepatic function [Increased AST (GOT) , ALT (GPT) etc]

Note 1) If such symptoms are observed, administration should be discontinued.

(3) Use in the Elderly

Because elderly patients often have reduced physiological function, careful supervision and measures such as reducing the dose are recommended.

(4) Use during Pregnancy, Delivery or Lactation

The safety of this product in pregnant women has not been established. Therefore, the product should be used in pregnant women, women who may possibly be pregnant only if the expected therapeutic benefits outweigh the possible risks associated with treatment.

(5) Pediatric Use

The safety of this product in children has not been established.
[Insufficient clinical data]

PRECAUTIONS FOR HANDLING

- Store at dry and cool place, protected from direct sunlight.
- Since this product contains natural crude drugs, some differences may be noted in the color or taste, etc. However, there is no change in the effect.

PACKAGING

168 g (2.0 g x 84 packets)

REQUEST FOR LITERATURE SHOULD BE MADE TO:

Dep. of PMS Information,
Ohsugi Pharmaceutical Co., Ltd.
1-8-6, Yamasaka, Higashiumiyoshi-ku, Osaka 546-0035
050-3776-0358