

October 2005  
(2nd version)Standard Commodity Classification No. of Japan  
875200

- Kampo product -

**JUNKOU Tokakujokito FC Extract Fine Granules for Ethical Use  
(Tokakujokito)**

Storage: Store at room temperature.  
See the section "PRECAUTIONS FOR HANDLING"  
Expiration date: The expiration date is indicated on the outer package.

Approval No.	(61AMY) 0419
Date of listing in the NHI reimbursement price	October 1986
Date of initial marketing in Japan	October 1986

**DESCRIPTION**

- (1) The daily dose of this product, 6.00 g, contains 2.30 g of the dried extract (Tokakujokito extract) from the following mixed crude drugs.
- |   |        |
|---|--------|
| JP Cinnamon Bark -----  | 4.00 g |
| Sodium sulfate Hydrate -----  | 2.00 g |
| JP Rhubarb -----  | 3.00 g |
| JP Peach Kernel -----   | 5.00 g |
| JP Glycyrrhiza -----  | 1.50 g |
| (JP: The Japanese Pharmacopeia)   |        |
| The inactive ingredients contained are Corn Starch and Lactose Hydrate. |        |
- (2) This product is light brown-colored fine granules, smells uniquely, and tastes bitter initially and slightly sweet later.
- ID Code: FC 61

**INDICATIONS**

The following symptoms of those patients with a comparatively strong constitution and hot flashes who are likely to have constipation:  
Menstrual irregularity, dysmenorrhea, anxiety during menstruation or following childbirth, low back pain, constipation, accessory symptoms associated with hypertension (headache, dizziness, and shoulder stiffness)

**DOSAGE AND ADMINISTRATION**

The usual adult dose is 6.0 g/day orally in 2 or 3 times before or between meals. The dosage may be adjusted according to the patient's age, body weight, and symptoms.

**PRECAUTIONS****(1) Careful Administration (Tokakujokito should be administered with care in the following patients.)**

- Patient with diarrhea or soft feces [These symptoms may be aggravated]
- Patients with an extremely weak gastrointestinal tract [Anorexia, epigastric distress, abdominal pain, diarrhea, etc. may occur.]
- Patients with an extremely declined constitution [Adverse reactions are likely to occur, and the symptoms may be aggravated.]

**(2) Important Precautions**

- When this product is used, the patient's "SHO" (constitution/symptoms) should be taken into account. The patient's progress should be carefully monitored, and if no improvement in symptoms/findings is observed, continuous treatment should be avoided.

- Since this product contains Glycyrrhiza, careful attention should be paid to the serum potassium level, blood pressure, etc., and if any abnormality is observed, administration should be discontinued.
- When this product is coadministered with other Kampo-products (Japanese traditional herbal medicines), etc., attention should be paid to the duplication of the contained crude drugs. Special caution should be exercised when this product is coadministered with other products containing Rhubarb.
- Since there is an individual difference in the cathartic action of Rhubarb, caution should be exercised concerning the dosage and administration.

**(3) Drug Interactions**

Precautions for coadministration (Tokakujokito should be administered with care when coadministered with the following drugs.)

Drugs	Signs, Symptoms, and Treatment	Mechanism and Risk Factors
(1) Preparations containing Glycyrrhiza	Pseudoaldosteronism is likely to occur. Besides, myopathy is likely to occur as a result of hypokalemia. (Refer to the section "Clinically significant adverse reactions")	Since glycyrrhizinic acid has an accelerating action on the potassium excretion at the renal tubules, an acceleration of decrease in the serum potassium level has been suggested.
(2) Preparations containing glycyrrhizinic acid or glycyrrhizinates		

**(4) Adverse Reactions**

This product has not been investigated (drug use investigations, etc.) to determine the incidence of adverse reactions. Therefore, the incidence of adverse reactions is not known.

**1) Clinically significant adverse reactions**

① **Pseudoaldosteronism:** Pseudoaldosteronism such as hypokalemia, increased blood pressure, retention of sodium/body fluid, edema, increased body weight, etc. may occur. The patient should be carefully monitored (measurement of serum potassium level, etc.) and if any abnormality is observed, administration should be discontinued and appropriate measures such as administration of potassium preparations should be taken.

② **Myopathy:** Myopathy may occur as a result of hypokalemia. The patient should be carefully monitored, and if any abnormality such as weakness, convulsion/paralysis of limbs, etc. are observed, administration should be discontinued and appropriate measures such as administration of potassium preparations should be taken.

**2) Other adverse reactions**

	<b>Incidence Unknown</b>
<b>Hypersensitivity</b> Note 1)	Rash, Redness, Pruritus, etc.
<b>Gastrointestinal</b>	Anorexia, Epigastric distress, Abdominal pain, Diarrhea, etc.

Note 1) If such symptoms are observed, administration should be discontinued.

**(5) Use in the Elderly**

Because elderly patients often have reduced physiological function, careful supervision and measures such as reducing the dose are recommended.

**(6) Use during Pregnancy, Delivery or Lactation**

- 1) Use of this product in pregnant women, women who may possibly be pregnant is not recommended. [Rhubarb (uterotonic action and congestive action on the intrapelvic organs), anhydrous Mirabilium (uterotonic action), and Peach Kernel contained in this product may cause premature birth or abortion.]
- 2) This product should be administered with care in nursing mothers. [Anthraquinone derivatives in Rhubarb contained in this product may be excreted in the breast milk and induce diarrhea in nursing infants.]

**(7) Pediatric Use**

The safety of this product in children has not been established.  
[Insufficient clinical data]

**(8) Other Precautions**

Since this product contains Mirabilium, caution should be exercised when continuous treatment with this product is given to patients who need limited salt-intake therapeutically.

**PRECAUTIONS FOR HANDLING**

- Store at dry and cool place, protected from direct sunlight.
- Since this product contains natural crude drugs, some differences may be noted in the color or taste, etc. However, there is no change in the effect.

**PACKAGING**

168 g (2.0 g x 84 packets)

**REQUEST FOR LITERATURE SHOULD BE MADE TO:**

Dep. Of PMS Information,  
Ohsugi Pharmaceutical Co., LTD.  
1-8-6, Yamasaka, Higashisumiyoshi-ku, Osaka 546-0035  
050-3776-0358