

- Kampo product -
Powerful Drug (Large Package only)
OHSUGI Hachimijioganryo Extract Granules G
(Hachimijiogan)

Storage: Store at room temperature.
See the section "PRECAUTIONS FOR HANDLING"
Expiration date: The expiration date is specified on the container or the outer package.

Approval No.	(61AM) 3901
Date of listing in the NHI reimbursement price	October 1986
Date of initial marketing in Japan	October 1986

DESCRIPTION

(1) The daily dose of this product, 7.5 g, contains 4.6 g of the dried extract (Hachimijioganryo extract) from the following mixed crude drugs.

JP Rehmannia Root -----	5 g
JP Poria Sclerotium -----	3 g
JP Cornus Fruit -----	3 g
JP Moutan Bark -----	3 g
JP Dioscorea Rhizome -----	3 g
JP Cinnamon Bark -----	1 g
JP Alisma Tuber -----	3 g
JP Powdered Processed Aconite Root -----	1 g

(JP: The Japanese Pharmacopeia)

The inactive ingredients contained are Lactose Hydrate, Corn Starch and Magnesium Stearate.

(2) This product is light grayish brown to light grayish and dark brown-colored granules, smells uniquely, and tastes slightly bitter and acidic.

ID Code: SG-07

INDICATIONS

The following symptoms of those patients with decreased urine volume or polyuria sometimes having dry mouth who are easily fatigued and easily feel cold in the extremities: Leg pain, low back pain, numbness, blurred vision in old patients, pruritus, dysuria, frequent urination, and edema

DOSAGE AND ADMINISTRATION

The usual adult dose is 7.5 g/day orally in 2 or 3 times before or between meals. The dosage may be adjusted according to the patient's age, body weight, and symptoms.

PRECAUTIONS

(1) Careful Administration (Hachimijioganryo should be administered with care in the following patients.)

- 1) Patients with strong constitution [Adverse reactions are likely to occur, and the symptoms may be aggravated.]
- 2) Patients with sensitivity to heat, a tendency towards hot flushes and red face.
- 3) Patients with an extremely weak gastrointestinal tract [Anorexia, epigastric distress, nausea, vomiting, abdominal pain, diarrhea, constipation etc. may occur.]
- 4) Patients with anorexia, nausea or vomiting [These symptoms may be aggravated]

(2) Important Precautions

- 1) When this product is used, the patient's "SHO" (constitution/symptoms) should be taken into account. The patient's progress should be carefully monitored, and if no improvement in symptoms/findings is observed, continuous treatment should be avoided.
- 2) When this product is coadministered with other Kampo-products (Japanese traditional herbal medicines), etc., attention should be paid to the duplication of the contained crude drugs. Special caution should be exercised when this product is coadministered with preparation containing Aconite Root.

(3) Adverse Reactions

This product has not been investigated (drug use investigations, etc.) to determine the incidence of adverse reactions. Therefore, the incidence of adverse reactions is not known.

	Incidence Unknown
Hypersensitivity Note 1)	Rash, Redness, Pruritus, etc.
Hepatic	Abnormality of hepatic function [increased AST (GOT), ALT (GPT), and total bilirubin levels, etc.]
Gastrointestinal	Anorexia, Epigastric distress, Nausea, Vomiting, Abdominal pain, Diarrhea, constipation, etc.
Other	Palpitation, Hot flush, Numbness of the tongue, etc.

Note 1) If such symptoms are observed, administration should be discontinued.

(4) Use in the Elderly

Because elderly patients often have reduced physiological function, careful supervision and measures such as reducing the dose are recommended.

(5) Use during Pregnancy, Delivery or Lactation

Use of this product in pregnant woman, women who may possibly be pregnant is not recommended. [Moutan Bark contained in this product may cause premature birth or abortion. Besides, adverse reactions due to Powdered Processed Aconite Root are likely to occur.]

(6) Pediatric Use

This product should be administered with care in children. [This product contains Powdered Processed Aconite Root.]

PRECAUTIONS FOR HANDLING

- Store at dry and cool place, protected from direct sunlight.
- Since this product contains natural crude drugs, some differences may be noted in the color or taste, etc. However, there is no change in the effect.

PACKAGING

500g
735g (2.5g x 294 packets)
210g (2.5g x 84 packets)

REQUEST FOR LITERATURE SHOULD BE MADE TO:

Dep. of PMS Information,
Ohsugi Pharmaceutical Co., Ltd.
1-8-6, Yamasaka, Higashisumiyoshi-ku, Osaka 546-0035
050-3776-0358