

- Kampo product -

**OHSUGI Mashininganryo Extract Granules G**  
(Mashiningan)

Storage: Store at room temperature.  
See the section "PRECAUTIONS FOR HANDLING"  
Expiration date: The expiration date is specified on the container or the outer package

Approval No.	(61AM) 3907
Date of listing in the NHI reimbursement price	October 1986
Date of initial marketing in Japan	October 1986

**DESCRIPTION**

(1) The daily dose of this product, 6.0g, contains 2.6g of the dried extract (Mashiningan extract) from the following mixed crude drugs.

JP Hemp Fruit -----	5 g
JP Magnolia Bark -----	2 g
JP Peony Root -----	2 g
JP Rhubarb -----	4 g
JP Immature Orange -----	2 g
JP Apricot Kernel -----	2 g

(JP: The Japanese Pharmacopeia)

The inactive ingredients contained are Lactose Hydrate, Corn Starch and Magnesium stearate.

(2) This product is light grayish dark brown-colored granules, smells uniquely, and tastes bitter and astringent.  
ID Code: SG-126

**INDICATIONS**

Constipation

**DOSAGE AND ADMINISTRATION**

The usual adult dose is 6.0 g/day orally in 2 or 3 times before or between meals. The dosage may be adjusted according to the patient's age, body weight, and symptoms.

**PRECAUTIONS****(1) Careful Administration (Mashiningan should be administered with care in the following patients.)**

- 1) Patient with diarrhea or soft feces [These symptoms may be aggravated]
- 2) Patients with an emarkably weak gastrointestinal tract [Anorexia, abdominal pain, diarrhea, etc. may occur.]

**(2) Important Precautions**

- 1) When this product is used, the patient's "SHO" (constitution/symptoms) should be taken into account. The patient's progress should be carefully monitored, and if no improvement in symptoms/findings is observed, continuous treatment should be avoided.
- 2) When this product is coadministered with other Kampo-products (Japanese traditional herbal medicines), etc., attention should be paid to any duplication of the contained crude drugs. Special caution should be exercised when this product is coadministered with other products containing Rhubarb.
- 3) Since there is an individual differenc in the cathartic action of Rhubarb, caution should be exercised concerning the dosage and administration.

**(3) Adverse Reactions**

This product has not been investigated (drug use investigations, etc.) to determine the incidence of adverse reactions. Therefore, the incidence of adverse reactions is not known.

	Incidence Unknown
--	-------------------

<b>Gastrointestinal</b>	Anorexia, Abdominal pain, Diarrhea, etc.
-------------------------	--

**(4) Use in the Elderly**

Because elderly patients often have reduced physiological function, careful supervision and measures such as reducing the dose are recommended.

**(5) Use during Pregnancy, Delivery or Lactation**

- 1) Use of this product in pregnant women, women who may possibly be pregnant is not recommended. [The uterotonic action and congestive action on the intrapelvic organs of Rhubarb contained in this product may cause premature birth or abortion.]
- 2) This product should be administered with care in nursing mothers. [Anthraquinone derivatives in Rhubarb contained in this product may be excreted in breast milk and induce diarrhea in nursing infants.]

**(6) Pediatric Use**

The safety of this product in children has not been established. [Insufficient clinical data]

**PRECAUTIONS FOR HANDLING**

- Store at dry and cool place, protected from direct sunlight.
- Since this product contains natural crude drugs, some differences may be noted in the color or taste, etc. However, there is no change in the effect.

**PACKAGING**

500g  
588g (2.0g x 294 packets)  
168g (2.0g x 84 packets)

**REQUEST FOR LITERATURE SHOULD BE MADE TO:**

Dep. of PMS Information,  
Ohsugi Pharmaceutical Co., LTD.  
1-8-6, Yamasaka, Higashisumiyoshi-ku, Osaka 546-0035  
050-3776-0358