

- Kampo product -

**OHSUGI Shireito Fine Granules for Ethical Use**  
(Shireito)

Storage: Store at room temperature.  
See the section "PRECAUTIONS FOR HANDLING"  
Expiration date: The expiration date is specified on the container or the outer package

Approval No.	(54AM) 1607
Date of listing in the NHI reimbursement price	September 1981
Date of initial marketing in Japan	September 1981

**DESCRIPTION**

- (1) The daily dose, 3.0g, of this product contains following powdered crude drugs.
- JP Powdered Alisma Tuber ----- 0.75 g  
JP Powdered Atractylodes Lancea Rhizome ----- 0.75 g  
JP Powdered Poria Sclerotium ----- 0.75 g  
JP Powdered Polyporus Sclerotium ----- 0.75 g  
(JP: The Japanese Pharmacopeia)
- (2) This product is dark brown to grayish brown-colored fine granules, smells uniquely, and tastes slightly bitter.  
ID Code: SG-140

**(3) Use during Pregnancy, Delivery or Lactation**

The safety of this product in pregnant woman has not been established. Therefore, the product should be used in pregnant women, women who may possibly be pregnant only if the expected therapeutic benefit outweigh the possible risks associated with treatment.

**(4) Pediatric Use**

The safety of this product in children has not been established.  
[Insufficient clinical data]

**INDICATIONS**

The following symptoms of those patients with thirst, decreased urine output despite water intake, and any of nausea, abdominal pain, edema, etc.:

Illness caused by the summer heat, acute gastroenteritis, and edema

**DOSAGE AND ADMINISTRATION**

The single dose is 1.0 g for adult. Two third of the adult dose for the children of the age from 7 years up and under 15years, a half of the adult dose for those of the age from 2 years up and under 7 years, and one forth of the adult dose for those of the age under 2 years. The single dose should be taken 3 times a day with water or hot water before or between meals.

**PRECAUTIONS****(1) Important Precautions**

- When this product is used, the patient's "SHO" (constitution/symptoms) should be taken into account. The patient's progress should be carefully monitored, and if no improvement in symptoms/findings is observed, continuous treatment should be avoided.
- When this product is coadministered with other Kampo-products (Japanese traditional herbal medicines), etc., attention should be paid to any duplication of the contained crude drugs.

**(2) Use in the Elderly**

Because elderly patients often have reduced physiological function, careful supervision and measures such as reducing the dose are recommended.

**PRECAUTIONS FOR HANDLING**

- Store at dry and cool place, protected from direct sunlight.
- Since this product contains natural crude drugs, some differences may be noted in the color or taste, etc. However, there is no change in the effect.

**PACKAGING**

500g  
294g (1.0g x 294 packets)  
84g (1.0g x 84 packets)

**REQUEST FOR LITERATURE SHOULD BE MADE TO:**

Dep. of PMS Information,  
Ohsugi Pharmaceutical Co., LTD.  
1-8-6, Yamasaka, Higashiumiyoshi-ku, Osaka 546-0035  
050-3776-0358